

# Zoonoses of Rural and Agricultural Occupations:

*Principles and common examples*

---


NE/CS-CASH. Ag Med Core Course  
July 13, 2022

- ❑ Kelley J. Donham MS, DVM, DACVPM
- ❑ Emeritus Professor, Agricultural Medicine, University of Iowa College of Public Health ([Kelley-Donham@uiowa.edu](mailto:Kelley-Donham@uiowa.edu))



# Bibliography

---



Donham and Thelin, 2016  
Wiley-Blackwell

Zoonoses - Infections Affecting Humans and Animals

Sing, A, Springer, 2015

Zoonoses: Recognition, Control, and Prevention

Hugh-Jones, M., Hubbert, W., Hagstad, H.

Wiley & Sons, 2008

Zoonosis emergence linked to agricultural intensification and environmental change  
[Bryony A. Jones](#), [Delia Grace](#), [Richard Kock](#), [Proc Natl Acad Sci U S A](#). 2013 May 21; 110(21): 8399–8404.

Human–livestock contacts and their relationship to transmission of zoonotic pathogens, a systematic review of literature  
Gijsklous, Heederik, Coutinho

One Health 2; pp 65-76


December 2016, Pages 65-76

open access

# Topics Covered

---



- Overview and general epidemiology
- Endemic and Epidemic Zoonoses
- Epidemic Zoonoses
  - Influenza A
  - Corona Virus
- Endemic Zoonoses
  - Leptospirosis
  - Erysipeloid
  - *Strep suis*
  - MRSA (Methicillin resistant *Staphylococcus aureus*)
  - Tetanus



# General Features of Agricultural Zoonoses

---

1. Zoonoses are diseases common to animals and man
2. Historically they have changed history.



2. There are over 250 zoonoses in the world
3. 60% Human pathogens, and 75% of emerging diseases = zoonotic. <http://fazd.tamu.edu/>

(Donham and Thelin, 2006, pp 357 - 380)

E.G.: Influenza, CORONA, HIV, ebola, Mad Cow, zika, nipa

---

# General Features of Agricultural Zoonoses

4. The risk of contracting a zoonotic disease depends on activities that bring humans into close association with animals/environment

5. Twenty-four of these are hazards for agricultural workers in the U.S.



# General Features of Agricultural Zoonoses

---

6. Agricultural zoonoses can be classified by their “relative risk” with type of livestock, e.g. swine, dairy, beef, poultry, or the general outdoor environment

# Zoonoses can be classified by their “relative risk” with type of livestock, e.g. swine, dairy, beef, poultry, or the general outdoor environment

## BEEF CATTLE:

Anthrax  
BSE  
Rabies  
Leptospirosis



## DAIRY CATTLE:

Milker's nodule  
Q Fever  
Staph infection  
Vesicular Stomatitis  
Zoophilic Ringworm



## POULTRY:

Histoplasmosis  
Newcastle disease  
Influenza  
Ornithosis



## SHEEP:

Contagious ecthyma  
Brucellosis  
Hydatid disease  
Tularemia



## SWINE:

Leptospirosis  
Erysipeloid  
Swine influenza  
NIPAH Virus  
S. suis  
Hepatitis E  
MRSA



## RURAL ENVIRONMENT:

Blastomycosis  
Arthropod-borne  
Viral encephalitis  
Rocky Mountain Spotted  
Tetanus  
Toxoplasmosis



# Epidemiologic “generalities” of Zoonoses

---

7. Seven general characteristics of these diseases:
  - a. They have non-specific symptoms, often resembling severe influenza
  - b. They are difficult to diagnose
  - c. They cause illness, but are rarely fatal
  - d. Animals are often sub-clinical chronic carriers
  - e. Humans are the dead end hosts
  - f. They cause economic losses when livestock are affected
  - g. Human cases are usually sporadic, (epidemics uncommon)



## Vulnerable Populations:

- Children
  - Lower immunity
- Those lacking “herd immunity”
- Women
  - Abortion risks
    - Brucella,
    - Q Fever,
    - Listeria


5/31/22



# Pathogenesis of Infectious Agents

---

## The Structure of Viruses



- ❑ “Germs” are parasites
- ❑ A battle between host defenses and the mechanisms of the agent.
- ❑ Example of viruses entering cells.



# Endemic and Epidemic Zoonoses

---

## □ Epidemic viruses

- Influenza viruses

- Corona Viruses



# Zoonotic Influenza

---

*Swine, Avian, Human  
Horses and Dogs*

(Donham and Thelin, 2006 p 371, Capua, 2013 [Vet Microbiol.](#) 2013  
26;165(1-2):7-12 )

# Influenza Virus

---

- Family Orthomyxoviridae

- “myxo” means mucus

- Three main types

- Type A

- Zoonotic strains

- Multiple species

- Type B


- Humans

- Type C


- Humans and swine



# Genetics of Influenza A



# Interspecies Transmission



RED ARROWS: Indicate transmission has occurred  
LIGHT ARROWS: Not reported, but show virus potential

## Viruses can transform into swine flu

Swine flu regularly causes outbreaks of influenza in pigs, but human infections have sporadically occurred.


### How swine flu spreads

Various species infected by the flu viruses ...

... can infect pigs, swapping genes, forming new viruses ...

... that can infect humans who have direct exposure to pigs ...

... who in turn can infect other humans.




SOURCE: Centers for Disease Control and Prevention

AP

# Annual influenza and periodic pandemics

---





# Zoonotic Influenza History and Biology

---



# The "Spanish Flu H1N1" 1918-1919

50 – 100 million estimated deaths



*I had a little bird  
Its name was Enza  
I opened a window  
And in-flu-enza*



# 1976 H1N1 Swine Flu Re-emerges

---



A/New Jersey/76? =

A/Swine [HSW<sub>1</sub>N<sub>1</sub>] =

Virus of 1918-1919




# 2009 Swine Flu Again? (Novel H1N1)



State of Veracruz  
First cases of  
H1N1 2009

# Now, Novel H1N1 of 2009 aka "Swine Flu"

---



# Avian Influenza in North America 2015 & 2022

(<http://www.cdc.gov/flu/avian/outbreaks/current.htm>)


(<https://www.cdc.gov/flu/avianflu/north-american-lineage.htm>)

- ❑ Most low path – no problem
- ❑ High Path (most H5 N1 something)
- ❑ Low transmission to humans
- ❑ One diagnosed Human illnesses in U.S >
- ❑ Asian & Arica human cases 447 fatalities
- ❑ [http://www.who.int/influenza/human\\_animal\\_interface/EN\\_GIP\\_20150501\\_CumulativeNumberH5N1cases.pdf?ua=1](http://www.who.int/influenza/human_animal_interface/EN_GIP_20150501_CumulativeNumberH5N1cases.pdf?ua=1)



North American Lineage avian viruses Low risk transmission to humans (H5N2 in 2015)

## H5N2 Influenza Spread, 2015




As of May 31. Source: USDA

# Bird Flu –Where did it come from where is it going?

- ❑ High Path Strain
- ❑ H5N1 mainly
- ❑ Outbreaks in 2015, and 2022
- ❑ Origan wild migrating waterfowl.
- ❑ Human not very susceptible (1 case in U.S. - no sx.)
- ❑ Asia & Africa since 2003. 864 cases 456 fatal)
- ❑ **Avian Influenza A (H5N1) - United States of America , WHO 5/6/22**




**H5N2 Influenza Spread, 2015**



# Corona Viruses

---

- ❑ RNA virus
- ❑ Surface binding proteins allow entry to cells
- ❑ RNA recombination results in Variation in pathologic and infectious capability
- ❑ Many different strains that can infect animals and humans





# History of Corona viruses infecting humans

- ❖ Many common Corona viruses
- ❖ 3 NOVEL corona viruses have resulted in epidemics

## SARS

- (severe Acute Respiratory syndrome)
- 2003 -2004
- Origin – China
- Bats -- Civet Cat –People
- Pandemic – but brief
- 8000 total human cases

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3747533/>


## MERS

- Middle East Respiratory Syndrome
- 2012 (still around)
- Origin – Saudi Peninsula (localized there)

## COVID 19 PANDEMIC

- July 2021. 1.9 million cases > 4million deaths. World wide. U.S. >34 million cases, >600 deaths.
- 188 countries

<https://coronavirus.jhu.edu/map.html>



# (SARS-CoV-2) COVID 19)



- ❑ February 2020 - ???
- ❑ China
- ❑ "Wet" markets
- ❑ Bats
- ❑ Pangolins?
- ❑ Laboratory Escape?
- ❑ Continued research

# Animal species susceptible to COVID 19

Domestic livestock/poultry

Pets

Other

Cows	Dogs	Bats
Pigs	Cats	Pangolins
Chickens	Ferrets	Mink
Ducks		Mice
		Lions
		Tigers
		Camels
		Primates



CDC: [https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/sa\\_one\\_health/sars-cov-2-animals-us](https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/sa_one_health/sars-cov-2-animals-us)

USDA: <https://www.cdc.gov/coronavirus/2019-ncov/daily-life-coping/animals.html>

# Low Risk of transmission of COVID 19 between animals and humans

From the CDC

- Re Pets

- "At this time, there is **no evidence** that (domestic) animals play a significant role in spreading the virus that causes COVID-19".
- "**People sick with COVID-19 should isolate themselves from other people and animals, including pets, during their illness until we know more about how this virus affects animals**".

- Re Livestock

- **No known risk of humans infected from pigs or cows.**
- **Risk to farmers is emotional and economic because of euthanizing pigs as slaughter plants closed because of COVID 19**



# Monkey Pox ??

- Not new
  - 2003 – import of African pet prairied dogs
    - 2022 – Human to human (sexual trans?)
- Trans by Direct contact, not aerosol like COVID
- Not a big threat
- Not an Agricultural risk



# Endemic Zoonoses

---



# Swine Zoonoses

---



5/31/22

# Case Study

---

- Malais
- 103° F
- Chills
- Muscular aches
  
- Headache
- Stiff neck
- Photophobia







5/31/22



5/31/22

# Leptospirosis

---

- *Leptospira interrogans*
- Worldwide distribution



5/31/22



5/31/22

# Treatment/Control and Prevention

---

- Antibiotics
  - Tetracycline
  - Penicillin
  - Streptomycin
  - Erythromycin
- Vaccination
- Caution in handling tissues
- Avoid direct contact with water/urine of potentially infected animals.



5/31/22



5/31/22





5/31/22

---

*Erysipelothrix rhusiopathiae*  
"Erysipelas" In pigs

# Pigs mainly but there are Other Sources of Erysipeloid

---

- ❑ Various Livestock species
- ❑ Soil
- ❑ Contaminated objects (fomites)

# Infection in People: Typically on Hand or Foot

- ❑ Swelling
- ❑ Deep burning, throbbing pain
- ❑ Skin tense
- ❑ No suppuration
- ❑ Violet-colored zone of erythema surrounding lesion
- ❑ Joints of phalanges, tender  
limited movement
- ❑ Axillary lymph nodes, swollen  
and tender
- ❑ Lesions on other body parts





Case #1 - Swine producer stopped by police - driving erratically

- ❑ No evidence of alcohol or drugs
- ❑ Taken to hospital
  - ❑ Severe septicemia
  - ❑ High fever, DIC
  - ❑ Meningitis
  - ❑ Lived, but with permanent CNS damage, extensive skin loss.
- ❑ *Streptococcus suis* was isolated

Case #2 – New York Farmer – Hospitalized for Meningitis – *S. suis* isolate, recently purchases piglets

# Streptococcus suis

---

- ❑ A common disease of swine
- ❑ Can cause infections in humans
- ❑ Septicemia, meningitis
- ❑ 40% of hospitalized cases = fatal
- ❑ Permanent brain damage – especially 8<sup>th</sup> cranial nerve function (hearing and balance)
- ❑ (<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2634616/>)
- ❑ [Emerg Infect Dis.](#) 2014 Jul; 20(7): 1105–1114.

# Streptococcus suis

❑ Nursery pigs most commonly affected – Meningitis

---


❑ Weak, unable to stand or walk

❑ Seizures

❑ Possible arthritis

❑ High mortality


❑ Sows are carriers - up to 80% of herd



# Primary Epidemiological Aspects of Streptococcus suis

## In the human population

- ❑ Mechanisms of transmission: contact with infected pigs or their environment; consuming contaminated pork.
- ❑ Population at risk: pork producers
- ❑ Recent China outbreak, 38 fatalities/215 cases
- ❑ 2008 sero-survey – 10% of swine exposed persons
- ❑ Misdiagnosed/under diagnosed/variance in virulence (Smith et. al. 2008)
  
- ❑ [Feng et.al. Virulence](#). 2014,5(4): 477–497.
- ❑ [Emerg Infect Dis](#). 2014 Jul; 20(7): 1105–1114.





# Control / Eradication:

---

## □ Good hygiene practices

- Environment – Power wash with biocide

- Personal

  - Wash hands

  - Treat Lacerations



## □ Keep *S. suis* out

- Biosecurity

- Test/treat/cull

- No Commercial Vaccine



# Joel - 1997

---



# MRSA


---

Methicillin-Resistant *Staphylococcus aureus*

(smith and Pearson 2010)

# Three Main Reservoirs of MRSA


---



Hospital (1980's)



Community  
(1990's)



Veal Calves - Netherlands (2004)  
Swine – Iowa (2009)

# Methicillin Resistant Staph aureus MRSA

---

- ❑ The “super bug”
- ❑ Since 1981 in the U.S.
- ❑ Initially hospital acquired. Now community acquired as well.
- ❑ Resistant to Methicillin and often several other beta lactams (penicillin group)
- ❑ Tetracycline resistance also common in swine LA MRSA strains
- ❑ 1% of the general population are carriers
- ❑ 15% - 40% of farm population LA MRSA carriers

# The Organism

---

## □ Staph aureus


- Gram positive coccus

## □ Virulence factors

- Adhere to surfaces
- Damage/avoid immune system

## □ Toxins

- Exotoxins – toxic shock, scalded skin
- Cytotoxin (PVL) – tissue necrosis
- Enterotoxins – preformed, gastroenteritis



# Epidemiology - MRSA

---

- Beta-lactam antibiotics damage bacterial cell wall & Penicillin binding sites (PBP)
  - MRSA Inactivates PBPs (penicillin. binding proteins)
- *mecA* gene codes for PBP<sub>a</sub>
  - Confers resistance to beta-lactam abx
  - Presence = Methicillin-resistance

[J Clin Microbiol](#). 2004 Dec; 42(12): 5881–5884

# MRSA Disease in Humans

---



# Clinical Signs – Humans

- Hospital-acquired
  - Wide variety of infections
  - Surgical site infections to invasive disease
- Community-acquired
  - Superficial skin, soft tissue disease
  - Pneumonia
  - Septicemia
  - Joint infections



# MRSA – Treatment

---

- Drainage and dressing
- Alternative to Beta Lactams antibiotics:
  - Trimethoprim/sulfamethoxazole (Bactrim)
  - Clindamycin
  - Gentamycin
  - Rifampin
- Decolonization – mupirocin (Bactroban)

# MRSA Disease in Animals

---

# MRSA in Animal Populations (this is a zoonotic infectious agent)

- Livestock: Pigs, Cattle, Horses
- ST 398 = Livestock Strain
- 60% of pigs infected mainly as carriers
- Mastitis in dairy cattle
- 15% - 40% of veterinarians colonized
- Pets: Dogs, cats
- Clinical disease rare



<http://img.photobucket.com/albums/312/halpe/mrsandog.jpg>



Smith, Livestock-Associated *Staphylococcus aureus*: The United States Experience

# Transmission

---

- Opportunistic in human
- Animals ↔ People ↔ Family
- Direct contact colonized people/animals
- Vertical spread from mother to fetus
- Fomites
  - Towels, used bandages
- Aerosol
- Oral (contaminated food)
- Clinical LA MRSA in U.S. unknown risk

# U. IA. Research on MRSA

Tara Smith PhD, Mike Male DVM, Dwight Ferguson, Abby Harper, Kerry Leedom DVM, MPH, Kelley Donham DVM

- ❑ Ecology?
- ❑ Pigs and people common/temporary carriers.
- ❑ Found in settled dust, air inside and outside swine buildings and shower facilities)
- ❑ Isolated from meat samples from grocery stores.
- ❑ Is it an important occupational or public health concern??



# Biosecurity issues?

---

- ✓ Where does it live in swine buildings?
- ✓ Its in animal feed
- ✓ Spread down wind
- ✓ How do we prevent its spread?
- ✓ Can we use bio-filters?



# Diagnosis

---

- Culture infection site
  - *Staph aureus* is coagulase positive
- Determine if *Staph aureus* is MRSA
  1. Antibiotic susceptibility testing
    - Oxacillin or cefoxitin
  2. Genetic testing
    - PCR to detect *mecA* gene
    - Livestock strains = St 398
    - Latex agglutination for PBP2a



# Prevention and Control

- ❑ Hygiene, hygiene, hygiene!!
- ❑ Cover skin abrasions  
Avoid sharing personal items
- ❑ Shower after exercising;  
clean equipment
- ❑ Screen health care &  
Swine workers
- ❑ Screen New Pts. in  
hospitals & nursing homes




# Summary of MRSA

---

- It is apparent that farm animals are a reservoir for MRSA
- There are new strains developing in the animal population
- Unknown occupational and public health risk.

# Tetanus

---



- Clostridium tetani
- Anaerobe – spore former – produces tetanospasmin
  - Blocks releasing factors of neurotransmitters at spinal cord level
- Lives in soil – feces from herbivore animals



# Tetanus Risk Factors

---


Anaerobic wounds contaminated with soil/herbivore feces

Deep puncture wounds

Tissue necrosis

Foreign body

Very young and elderly (insufficient immunity)



Infants of mothers not immunized or not breast feed

Bowel surgery

Contaminated needles



# Tetanus the Disease

---



# Tetanus Primary Prevention


---

- Tetanus Toxoid immunization
  - DPT Children (5 doses)
    - Spaced from 2 mo. – Beginning school age
  - Adults
    - 10 years
    - Or after severe exposure if > 5 years
    - Some recommend DTP in adults for booster

# A Quick Review

---

- Overview and general epidemiology
- Epidemic and Endemic conditions
- Leptospirosis
- Erysipeloid
- *Strep suis*
- MRSA (Methicillin resistant *Staphylococcus aureus*)
- Zoonotic Influenza
- Tetanus



# Post Quiz

---





Animals are the primary host for all Zoonoses communicable to man.

---

True

False

Don't know

Generally, a zoonotic infection in a person is readily transmitted from person to person.

---

True

False

Don't know

## Which of the following is NOT True?

---

- Zoonoses in humans usually occur in broad epidemics
- There are over 250 zoonoses in the world
- Zoonoses make up 60% of human pathogens and 75% of emerging diseases

Leptospirosis is disseminated by contact with urine of an infected pig, cow, raccoon, squirrel, or mouse.

---

True

False

Don't know

Methicillin resistant *Staphylococcus aureus* (MRSA) is a commonly recognized occupational disease of pork producers.

---

True

False

Don't know

# Which of the following is the greater risk for tetanus?

---

- Laceration on the hand while castrating pigs
  - Dairy farmer sticks his foot with a pitchfork while cleaning the barn
  - Stepping on a rusty nail
  - Don't know
-

# Human beings cannot acquire animal ring worm infection

---

True

False

Don't know